

周産期死亡

〔月別にみた周産期死亡〕
〔Perinatal deaths by months〕

表 8.3 月別にみた妊娠満28週以後の死産—早期新生児死亡・
年次別周産期死亡数及び率¹⁾ (出生千対)

Table 8.3 Trends of perinatal deaths and perinatal death rates by months : Japan

年次 Year	総数 Total	1月 Jan.	2月 Feb.	3月 Mar.	4月 Apr.	5月 May	6月 June	7月 July	8月 Aug.	9月 Sept.	10月 Oct.	11月 Nov.	12月 Dec.
周産期死亡数 Perinatal deaths													
1970 昭和45年	41 917	3 864	3 545	3 856	3 522	3 468	3 428	3 699	3 341	3 220	3 256	3 116	3 602
75 50	30 513	2 733	2 523	2 705	2 538	2 656	2 545	2 760	2 628	2 330	2 366	2 217	2 512
80 55	18 385	1 544	1 472	1 546	1 560	1 620	1 514	1 633	1 604	1 517	1 449	1 431	1 495
85 60	11 470	961	952	983	916	1 043	969	1 000	1 007	906	953	861	919
87 62	9 317	837	723	798	788	869	822	814	788	756	728	653	741
88 63	8 508	760	656	692	704	731	747	754	747	655	696	690	676
妊娠満28週以後の死産数 Foetal deaths of 28 weeks and over of gestation													
1970 昭和45年	29 107	2 766	2 522	2 746	2 497	2 437	2 385	2 514	2 261	2 209	2 207	2 105	2 458
75 50	20 268	1 860	1 711	1 858	1 706	1 769	1 659	1 748	1 764	1 528	1 549	1 442	1 674
80 55	12 231	1 054	991	1 027	1 069	1 080	988	1 105	1 073	985	986	921	952
85 60	7 733	659	654	656	630	671	644	690	682	609	626	607	605
87 62	6 252	560	481	550	546	579	560	543	519	498	489	428	499
88 63	5 759	525	453	478	471	501	506	510	498	435	454	467	461
早期新生児死亡数 Early neonatal deaths													
1970 昭和45年	12 810	1 098	1 023	1 110	1 025	1 031	1 043	1 185	1 080	1 011	1 049	1 011	1 144
75 50	10 245	873	812	847	832	887	886	1 012	864	802	817	775	838
80 55	6 154	490	481	519	491	540	526	528	531	532	463	510	543
85 60	3 737	302	298	327	286	372	325	310	325	297	327	254	314
87 62	3 065	277	242	248	242	290	262	271	269	258	239	225	242
88 63	2 749	235	203	214	233	230	241	244	249	220	242	223	215
周産期死亡率 Perinatal death rates													
1970 昭和45年	21.7	22.1	22.9	23.5	21.2	21.6	22.4	21.8	20.4	20.5	21.3	20.9	21.5
75 50	16.0	16.3	16.5	16.9	15.9	16.5	16.7	16.4	16.0	14.6	15.3	15.5	16.0
80 55	11.7	11.4	11.8	11.9	12.2	12.1	11.8	11.8	11.6	11.1	10.9	11.9	11.7
85 60	8.0	8.0	8.6	8.5	8.0	8.6	8.3	7.7	8.0	7.4	7.7	7.7	7.7
87 62	6.9	7.4	7.1	7.1	7.2	7.5	7.3	6.7	6.7	6.5	6.5	6.2	6.7
88 63	6.5	7.1	6.6	6.5	6.7	6.7	6.9	6.5	6.3	5.7	6.1	6.5	6.3
妊娠満28週以後の死産比 Foetal death ratio of 28 weeks and over of gestation													
1970 昭和45年	15.0	15.8	16.3	16.7	15.0	15.1	15.6	14.8	13.8	14.1	14.4	14.1	14.7
75 50	10.7	11.1	11.2	11.6	10.7	11.0	10.9	10.4	10.7	9.6	10.0	10.1	10.7
80 55	7.8	7.8	7.9	7.9	8.3	8.0	7.7	8.0	7.8	7.2	7.4	7.6	7.5
85 60	5.4	5.5	5.9	5.7	5.5	5.5	5.5	5.3	5.4	5.0	5.1	5.4	5.1
87 62	4.6	5.0	4.7	4.9	5.0	5.0	5.0	4.5	4.4	4.4	4.4	4.0	4.5
88 63	4.4	4.9	4.6	4.5	4.5	4.6	4.7	4.4	4.2	3.8	4.0	4.4	4.3
早期新生児死亡率 Early neonatal death rates													
1970 昭和45年	6.6	6.3	6.6	6.8	6.2	6.4	6.8	7.0	6.6	6.4	6.9	6.8	6.8
75 50	5.4	5.2	5.3	5.3	5.2	5.5	5.8	6.0	5.3	5.0	5.3	5.4	5.3
80 55	3.9	3.6	3.8	4.0	3.8	4.0	4.1	3.8	3.8	3.9	3.5	4.2	4.3
85 60	2.6	2.5	2.7	2.8	2.5	3.1	2.8	2.4	2.6	2.4	2.7	2.3	2.6
87 62	2.3	2.4	2.4	2.2	2.2	2.5	2.3	2.2	2.3	2.3	2.1	2.1	2.2
88 63	2.1	2.2	2.1	2.0	2.2	2.1	2.2	2.1	2.1	1.9	2.1	2.1	2.0

注：1) 計算方法は2ページの「月間周産期死亡率」を参照されたい。

Perinatal deaths

〔出産時の体重別にみた周産期死亡〕
〔Perinatal deaths by birth weight〕

表 8.4 性・体重別にみた妊娠満28週以後の死産—早期新生児死亡別周産期死
亡数・率(性・体重別出生千対) 及び百分率

Table 8.4 Perinatal deaths, perinatal death rates (per 1,000 live births)
and percent distribution by sex and birth weight : Japan, 1988

昭和63年

		総数 Total	～0.9kg	1.0～1.4	1.5～1.9	2.0～2.4	2.5～2.9	3.0～3.4	3.5～3.9	4.0～	(再掲) 2.5kg 以下 2.5kg and under (Secondary mention)
周産期死亡数 ¹⁾ Perinatal deaths											
総数 Total	男 M.	8 508	1 502	1 456	1 292	1 326	1 420	968	329	142	5 608
	女 F.	4 496	759	782	674	688	712	545	208	96	2 922
妊娠満28週以後の死産 Foetal death of 28 weeks and over of gestation	男 M.	5 759	862	1 112	970	946	956	604	199	97	3 908
	女 F.	3 006	418	580	512	487	485	323	131	66	2 008
早期新生児死亡 Early neonatal deaths	男 M.	2 749	640	344	322	380	464	364	130	45	1 700
	女 F.	1 490	341	202	162	201	227	222	77	30	914
周産期死亡率 ²⁾ Perinatal death rates											
総数 Total	男 M.	6.5	693.8	336.2	119.2	22.7	3.7	1.6	1.5	5.0	71.7
	女 F.	6.7	697.0	351.0	121.4	25.8	4.1	1.7	1.6	5.2	79.9
妊娠満28週以後の死産比 Foetal death ratio of 28 weeks and over of gestation	男 M.	6.1	645.9	313.8	115.0	19.8	3.4	1.5	1.4	4.3	62.6
	女 F.	4.4	398.2	256.8	89.5	16.2	2.5	1.0	0.9	3.4	49.9
早期新生児死亡率 Early neonatal death rates	男 M.	4.5	383.8	260.3	92.2	18.3	2.8	1.0	1.0	3.6	54.9
	女 F.	4.2	368.0	246.3	84.7	14.2	2.2	1.0	0.8	2.9	43.7
百分率 ³⁾ Percentage											
総数 Total	男 M.	100.0	17.8	17.3	15.3	15.7	16.8	11.5	3.9	1.7	66.5
	女 F.	100.0	17.0	17.5	15.1	15.4	15.9	12.2	4.7	2.2	65.5
妊娠満28週以後の死産 Foetal deaths of 28 weeks and over of gestation	男 M.	100.0	17.9	17.0	15.6	16.2	18.1	10.9	3.1	1.2	67.1
	女 F.	100.0	15.0	19.4	16.9	16.5	16.6	10.5	3.5	1.7	68.0
早期新生児死亡 Early neonatal deaths	男 M.	100.0	13.9	19.3	17.1	16.2	16.2	10.8	4.4	2.2	66.9
	女 F.	100.0	14.9	19.5	16.9	17.0	17.6	10.5	2.6	1.1	68.5
早期新生児死亡 Early neonatal deaths	男 M.	100.0	23.8	12.8	12.0	14.1	17.3	13.5	4.8	1.7	63.2
	女 F.	100.0	23.3	13.8	11.1	13.7	15.5	15.2	5.3	2.1	62.5
早期新生児死亡 Early neonatal deaths	男 M.	100.0	24.4	11.6	13.0	14.6	19.3	11.6	4.3	1.2	64.1
	女 F.	100.0	24.4	11.6	13.0	14.6	19.3	11.6	4.3	1.2	64.1

注：1) 性・体重ともに不詳を含む。
2) 周産期死亡率²⁾ = $\frac{\text{年間の性・体重階級別周産期死亡数}}{\text{年間の性・体重階級別出生数}} \times 1,000$
3) 体重不詳を除き算出した。