

周産期死亡

(月別にみた周産期死亡)  
(Perinatal deaths by months)

表 8.3 月別にみた妊娠満28週以後の死産—早期新生児死亡・  
年次別周産期死亡数及び率<sup>1)</sup> (出生千対)

Table 8.3 Trends of perinatal deaths and perinatal death rates by months : Japan

年次 Year	総数 Total	1月 Jan.	2月 Feb.	3月 Mar.	4月 Apr.	5月 May	6月 June	7月 July	8月 Aug.	9月 Sept.	10月 Oct.	11月 Nov.	12月 Dec.
<b>周産期死亡数 Perinatal deaths</b>													
1970 昭和45年	41 917	3 864	3 545	3 856	3 522	3 468	3 428	3 699	3 341	3 220	3 256	3 116	3 602
75 50	30 513	2 733	2 523	2 705	2 538	2 656	2 545	2 760	2 628	2 330	2 366	2 217	2 512
80 55	18 385	1 544	1 472	1 546	1 560	1 620	1 514	1 633	1 604	1 517	1 449	1 431	1 495
85 60	11 470	961	952	983	916	1 043	969	1 000	1 007	906	953	861	919
86 61	10 148	856	787	888	836	895	840	907	861	806	843	789	840
87 62	9 317	837	723	798	788	869	822	814	788	756	728	653	741
<b>妊娠満28週以後の死産数 Foetal deaths of 28 weeks and over of gestation</b>													
1970 昭和45年	29 107	2 766	2 522	2 746	2 497	2 437	2 385	2 514	2 261	2 209	2 207	2 105	2 458
75 50	20 268	1 860	1 711	1 858	1 706	1 769	1 659	1 748	1 764	1 528	1 549	1 442	1 674
80 55	12 231	1 054	991	1 027	1 069	1 080	988	1 105	1 073	985	986	921	952
85 60	7 733	659	654	656	630	671	644	690	682	609	626	607	605
86 61	6 902	586	554	607	546	611	564	624	588	559	593	504	566
87 62	6 252	560	481	550	546	579	560	543	519	498	489	428	499
<b>早期新生児死亡数 Early neonatal deaths</b>													
1970 昭和45年	12 810	1 098	1 023	1 110	1 025	1 031	1 043	1 185	1 080	1 011	1 049	1 011	1 144
75 50	10 245	873	812	847	832	887	886	1 012	864	802	817	775	838
80 55	6 154	490	481	519	491	540	526	528	531	532	463	510	543
85 60	3 737	302	298	327	286	372	325	310	325	297	327	254	314
86 61	3 246	270	233	281	290	284	276	283	273	247	250	285	274
87 62	3 065	277	242	248	242	290	262	271	269	258	239	225	242
<b>周産期死亡率 Perinatal death rates</b>													
1970 昭和45年	21.7	22.1	22.9	23.5	21.2	21.6	22.4	21.8	20.4	20.5	21.3	20.9	21.5
75 50	16.0	16.3	16.5	16.9	15.9	16.5	16.7	16.4	16.0	14.6	15.3	15.5	16.0
80 55	11.7	11.4	11.8	11.9	12.2	12.1	11.8	11.8	11.6	11.1	10.9	11.9	11.7
85 60	8.0	8.0	8.6	8.5	8.0	8.6	8.3	7.7	8.0	7.4	7.7	7.7	7.7
86 61	7.3	7.4	7.5	7.8	7.6	7.7	7.4	7.3	7.2	6.9	7.0	7.2	7.2
87 62	6.9	7.4	7.1	7.1	7.2	7.5	7.3	6.7	6.7	6.7	6.5	6.2	6.7
<b>妊娠満28週以後の死産比 Foetal death ratio of 28 weeks and over of gestation</b>													
1970 昭和45年	15.0	15.8	16.3	16.7	15.0	15.1	15.6	14.8	13.8	14.1	14.4	14.1	14.7
75 50	10.7	11.1	11.2	11.6	10.7	11.0	10.9	10.4	10.7	9.6	10.0	10.1	10.7
80 55	7.8	7.8	7.9	7.9	8.3	8.0	7.7	8.0	7.8	7.2	7.4	7.6	7.5
85 60	5.4	5.5	5.9	5.7	5.5	5.5	5.5	5.3	5.4	5.0	5.1	5.4	5.1
86 61	5.0	5.1	5.3	5.3	4.9	5.3	4.9	5.0	4.9	4.8	5.0	4.6	4.9
87 62	4.6	5.0	4.7	4.9	5.0	5.0	4.5	4.4	4.4	4.4	4.4	4.0	4.5
<b>早期新生児死亡率 Early neonatal death rates</b>													
1970 昭和45年	6.6	6.3	6.6	6.8	6.2	6.4	6.8	7.0	6.6	6.4	6.9	6.8	6.8
75 50	5.4	5.2	5.3	5.3	5.2	5.5	5.8	6.0	5.3	5.0	5.3	5.4	5.3
80 55	3.9	3.6	3.8	4.0	3.8	4.0	4.1	3.8	3.8	3.9	3.5	4.2	4.3
85 60	2.6	2.5	2.7	2.8	2.5	3.1	2.8	2.4	2.6	2.4	2.7	2.3	2.6
86 61	2.3	2.3	2.2	2.5	2.6	2.4	2.4	2.3	2.3	2.1	2.1	2.6	2.4
87 62	2.3	2.4	2.4	2.2	2.2	2.5	2.3	2.2	2.3	2.1	2.1	2.1	2.2

注：1) 計算方法は52ページの「月間周産期死亡率」を参照されたい。

Perinatal deaths

(出産時の体重別にみた周産期死亡)  
(Perinatal deaths by birth weight)

表 8.4 性・体重別にみた妊娠満28週以後の死産—早期新生児死亡別周産期死亡数・率(性・体重別出生千対)及び百分率

Table 8.4 Perinatal deaths, perinatal death rates (per 1,000 live births) and percent distribution by sex and birth weight : Japan, 1987

昭和62年

	総数 Total	1.0kg 未満 1.0kg	1.0~1.5	1.5~2.0	2.0~2.5	2.5~3.0	3.0~3.5	3.5~4.0	4.0kg 以上 4.0kg and over	(再掲) 2.5kg 以下 2.5kg and under (Secondary mention)
		<b>周産期死亡数<sup>1)</sup> Perinatal deaths</b>								
総数 Total	9 317	1 713	1 619	1 466	1 366	1 439	1 087	381	168	6 204
男 M.	5 018	891	881	744	737	781	599	236	109	3 274
女 F.	4 299	822	738	722	629	658	488	145	58	2 930
妊娠満28週以後の死産 Foetal death of 28 weeks and over of gestation	6 252	1 000	1 188	1 082	950	987	677	239	120	4 251
男 M.	3 292	487	625	559	499	525	371	149	74	2 186
女 F.	2 960	513	563	523	451	462	306	90	45	2 065
早期新生児死亡 Early neonatal deaths	3 065	713	431	384	416	452	410	142	48	1 953
男 M.	1 726	404	256	185	238	256	228	87	35	1 088
女 F.	1 339	309	175	199	178	196	182	55	13	865
<b>周産期死亡率<sup>2)</sup> Perinatal death rates</b>										
総数 Total	6.9	793.8	388.5	134.6	23.2	3.8	1.7	1.7	5.4	78.5
男 M.	7.2	810.0	399.7	135.6	27.2	4.5	1.8	1.7	5.6	88.2
女 F.	6.4	723.1	368.8	131.9	19.6	3.1	1.6	1.5	5.1	68.0
妊娠満28週以後の死産比 Foetal death ratio of 28 weeks and over of gestation	4.6	463.4	285.1	99.4	16.1	2.6	1.1	1.0	3.9	53.8
男 M.	4.8	442.7	283.6	101.9	18.4	3.0	1.1	1.1	3.8	58.9
女 F.	4.4	431.0	279.7	95.1	14.1	2.2	1.0	1.0	3.9	47.3
早期新生児死亡率 Early neonatal death rates	2.3	330.4	103.4	35.3	7.1	1.2	0.7	0.6	1.5	24.7
男 M.	2.5	367.3	116.2	33.7	8.8	1.5	0.7	0.6	1.8	29.3
女 F.	2.0	292.1	89.1	36.8	5.6	0.9	0.6	0.6	1.1	20.6
<b>百分率<sup>3)</sup> Percentage</b>										
総数 Total	100.0	18.5	17.5	15.9	14.8	15.6	11.8	4.1	1.8	67.2
男 M.	100.0	17.9	17.7	14.9	14.8	15.7	12.0	4.7	2.2	65.8
女 F.	100.0	18.3	17.4	17.1	15.0	15.7	11.7	3.5	1.4	68.2
妊娠満28週以後の死産 Foetal deaths of 28 weeks and over of gestation	100.0	16.0	19.0	17.3	15.2	15.8	10.8	3.8	1.9	68.1
男 M.	100.0	14.8	19.0	17.0	15.2	16.0	11.3	4.5	2.2	66.5
女 F.	100.0	15.9	19.2	17.9	15.6	16.0	10.6	3.1	1.6	69.2
早期新生児死亡 Early neonatal deaths	100.0	23.8	14.4	12.8	13.9	15.1	13.7	4.7	1.6	65.2
男 M.	100.0	23.9	15.2	11.0	14.1	15.2	13.5	5.2	2.1	64.4
女 F.	100.0	23.6	13.4	15.2	13.6	15.0	13.9	4.2	1.0	66.2

注：1) 性・体重ともに不詳を含む。  
2) 周産期死亡率 = 年間の性・体重階級別周産期死亡数 ÷ 年間の性・体重階級別出生数 × 1 000  
3) 体重不詳を除き算出した。