

周産期死亡

[月別にみた周産期死亡]
[Perinatal deaths by months]

表 8.3 月別にみた妊娠満28週以後の死産—早期新生児死亡・年次別周産期死亡数及び率¹⁾ (出生千対)

Table 8.3 Trends of perinatal deaths and perinatal death rates by months : Japan

年次	総数	1月	2月	3月	4月	5月	6月	7月	8月	9月	10月	11月	12月
Year	Total	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
周産期死亡数 Perinatal deaths													
1970	41 917	3 864	3 545	3 856	3 522	3 468	3 428	3 699	3 341	3 220	3 256	3 116	3 602
75	30 513	2 733	2 523	2 705	2 538	2 656	2 545	2 760	2 628	2 330	2 366	2 217	2 512
80	18 385	1 544	1 472	1 546	1 560	1 620	1 514	1 633	1 604	1 517	1 449	1 431	1 495
83	14 035	1 224	1 144	1 219	1 207	1 231	1 125	1 276	1 150	1 160	1 143	987	1 169
84	12 998	1 117	1 067	1 089	1 075	1 157	1 082	1 185	1 142	1 044	1 055	952	1 033
85	11 470	961	952	983	916	1 043	969	1 000	1 007	906	953	861	919
妊娠満28週以後の死産数 Late fetal deaths													
1970	29 107	2 766	2 522	2 746	2 497	2 437	2 385	2 514	2 261	2 209	2 207	2 105	2 458
75	20 268	1 860	1 711	1 858	1 706	1 769	1 659	1 748	1 764	1 528	1 549	1 442	1 674
80	12 231	1 054	991	1 027	1 069	1 080	988	1 105	1 073	985	986	921	952
83	9 464	825	769	849	849	799	757	863	767	775	784	663	764
84	8 724	764	704	725	731	781	752	802	736	681	713	636	699
85	7 733	659	654	656	630	671	644	690	682	609	626	607	605
早期新生児死亡数 Deaths under 1 week													
1970	12 810	1 098	1 023	1 110	1 025	1 031	1 043	1 185	1 080	1 011	1 049	1 011	1 144
75	10 245	873	812	847	832	887	886	1 012	864	802	817	775	838
80	6 154	490	481	519	491	540	526	528	531	532	463	510	543
83	4 571	399	375	370	358	432	368	413	383	385	359	324	405
84	4 274	353	363	364	344	376	330	383	406	363	342	316	334
85	3 737	302	298	327	286	372	325	310	325	297	327	254	314
周産期死亡率 Perinatal death rates													
1970	21.7	22.1	22.9	23.5	21.2	21.6	22.4	21.8	20.4	20.5	21.3	20.9	21.5
75	16.0	16.3	16.5	16.9	15.9	16.5	16.7	16.4	16.0	14.6	15.3	15.5	16.0
80	11.7	11.4	11.8	11.9	12.2	12.1	11.8	11.8	11.6	11.1	10.9	11.9	11.7
83	9.3	9.7	10.1	9.8	9.7	9.6	9.1	9.5	8.6	9.0	9.1	8.3	9.2
84	8.7	9.0	9.2	8.8	9.1	9.3	9.1	8.8	8.5	8.2	8.4	8.1	8.4
85	8.0	8.0	8.6	8.5	8.0	8.6	8.3	7.7	8.0	7.4	7.7	7.7	7.7
妊娠満28週以後の死産比 Late fetal death ratio													
1970	15.0	15.8	16.3	16.7	15.0	15.1	15.6	14.8	13.8	14.1	14.4	14.1	14.7
75	10.7	11.1	11.2	11.6	10.7	11.0	10.9	10.4	10.7	9.6	10.0	10.1	10.7
80	7.8	7.8	7.9	7.9	8.3	8.0	7.7	8.0	7.8	7.2	7.4	7.6	7.5
83	6.3	6.6	6.8	6.8	6.8	6.2	6.2	6.4	5.7	6.0	6.3	5.6	6.0
84	5.9	6.1	6.1	5.9	6.2	6.3	6.3	6.0	5.5	5.4	5.7	5.4	5.7
85	5.4	5.5	5.9	5.7	5.5	5.5	5.5	5.3	5.4	5.0	5.1	5.4	5.1
早期新生児死亡率 Death rates under 1 week													
1970	6.6	6.3	6.6	6.8	6.2	6.4	6.8	7.0	6.6	6.4	6.9	6.8	6.8
75	5.4	5.2	5.3	5.3	5.2	5.5	5.8	6.0	5.3	5.0	5.3	5.4	5.3
80	3.9	3.6	3.8	4.0	3.8	4.0	4.1	3.8	3.8	3.9	3.5	4.2	4.3
83	3.0	3.2	3.3	3.0	2.9	3.4	3.0	3.1	2.9	3.0	2.9	2.7	3.2
84	2.9	2.8	3.1	2.9	2.9	3.0	2.8	2.8	3.0	2.9	2.7	2.7	2.7
85	2.6	2.5	2.7	2.8	2.5	3.1	2.8	2.4	2.6	2.4	2.7	2.3	2.6

注： 1) 計算方法は52ページの「月間周産期死亡率」を参照されたい。

Perinatal deaths

[出産時の体重別にみた周産期死亡]
[Perinatal deaths by birth weight]

表 8.4 性・体重別にみた妊娠満28週以後の死産—早期新生児死亡別周産期死亡数・率 (性・体重別出生千対) 及び百分率

Table 8.4 perinatal deaths, perinatal death rates (per 1,000 live births) and percent distribution by sex and birth weight : Japan, 1985

		総数	1.0kg未満	1.0~1.5	1.5~2.0	2.0~2.5	2.5~3.0	3.0~3.5	3.5~4.0	4.0kg以上	(再掲) 2.5kg以下
		Total	Under 1.0kg	1.0~1.5	1.5~2.0	2.0~2.5	2.5~3.0	3.0~3.5	3.5~4.0	4.0kg and over	2.5kg and under (Secondary mention)
周産期死亡数¹⁾ Perinatal deaths											
総数	Total	1 1470	2 035	2 022	1 863	1 667	1 738	1 379	511	154	7 659
	男 M.	6 059	970	1 064	1 004	905	907	751	314	93	3 973
	女 F.	5 299	1 016	939	844	752	824	626	195	61	3 593
妊娠満28週以後の死産	Late fetal deaths	7 733	1 145	1 475	1 402	1 199	1 179	902	309	102	5 275
	男 M.	3 992	517	753	736	636	606	492	183	65	2 665
	女 F.	3 629	579	703	651	553	566	408	124	37	2 517
早期新生児死亡	Deaths under 1 week	3 737	890	547	461	468	559	477	202	52	2 384
	男 M.	2 067	453	311	268	269	301	259	131	28	1 308
	女 F.	1 670	437	236	193	199	258	218	71	24	1 076
周産期死亡率²⁾ Perinatal death rates											
総数	Total	8.0	944.8	435.3	162.6	27.8	4.4	2.1	2.0	4.2	93.2
	男 M.	8.2	949.1	438.2	173.3	33.0	5.1	2.2	2.1	4.1	103.3
	女 F.	7.6	897.5	423.5	149.0	23.2	3.8	2.0	1.8	4.5	82.2
妊娠満28週以後の死産比	Late fetal death ratio	5.4	531.6	317.5	122.4	20.0	3.0	1.4	1.2	2.8	64.2
	男 M.	5.4	505.9	310.1	127.0	23.2	3.4	1.4	1.2	2.8	69.3
	女 F.	5.2	511.5	317.1	115.0	17.0	2.6	1.3	1.2	2.7	57.6
早期新生児死亡率	Death rates under 1 week	2.6	413.2	117.8	40.2	7.8	1.4	0.7	0.8	1.4	29.0
	男 M.	2.8	443.2	128.1	46.3	9.8	1.7	0.7	0.9	1.2	34.0
	女 F.	2.4	386.0	106.5	34.1	6.1	1.2	0.7	0.7	1.8	24.6
百分率³⁾ Percentage											
総数	Total	100.0	17.9	17.8	16.4	14.7	15.3	12.1	4.5	1.4	67.4
	男 M.	100.0	16.1	17.7	16.7	15.1	15.1	12.5	5.2	1.5	66.1
	女 F.	100.0	19.3	17.9	16.1	14.3	15.7	11.9	3.7	1.2	68.3
妊娠満28週以後の死産	Late fetal deaths	100.0	14.8	19.1	18.2	15.5	15.3	11.7	4.0	1.3	68.4
	男 M.	100.0	13.0	18.9	18.5	15.9	15.2	12.3	4.6	1.6	66.8
	女 F.	100.0	16.0	19.4	18.0	15.3	15.6	11.3	3.4	1.0	69.5
早期新生児死亡	Deaths under 1 week	100.0	24.3	15.0	12.6	12.8	15.3	13.0	5.5	1.4	65.2
	男 M.	100.0	22.4	15.4	13.3	13.3	14.9	12.8	6.5	1.4	64.8
	女 F.	100.0	26.7	14.4	11.8	12.2	15.8	13.3	4.3	1.5	65.8

注： 1) 性・体重ともに不詳を含む。
2) 周産期死亡率 = $\frac{\text{年間の性・体重階級別周産期死亡数}}{\text{年間の性・体重階級別出生数}} \times 1000$
3) 体重不詳を除き算出した。