

周産期死亡

[月別にみた周産期死亡]

[Perinatal deaths by months]

表 8.3 月別にみた妊娠満28週以後の死産—早期新生児死亡・
年次別周産期死亡数及び率¹⁾ (出生千対)

Table 8.3 Trends of perinatal deaths and perinatal death rates
(per 1,000 live births) by months : Japan

年次	総数	1月	2月	3月	4月	5月	6月	7月	8月	9月	10月	11月	12月
Year	Total	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
周産期死亡数 Perinatal deaths													
1970 昭和45年	41 917	3 864	3 545	3 856	3 522	3 468	3 428	3 699	3 341	3 220	3 256	3 116	3 602
75 50	30 513	2 733	2 523	2 705	2 538	2 656	2 545	2 760	2 628	2 330	2 366	2 217	2 512
80 55	18 385	1 544	1 472	1 546	1 560	1 620	1 514	1 633	1 604	1 517	1 449	1 431	1 495
85 60	11 470	961	952	983	916	1 043	969	1 000	1 007	906	953	861	919
89 平成1年	7 450	666	524	621	626	614	620	633	642	635	626	619	624
90 2	7 001	636	532	625	617	582	577	558	613	569	581	551	560
妊娠満28週以後の死産数 Foetal deaths of 28 weeks and over of gestation													
1970 昭和45年	29 107	2 766	2 522	2 746	2 497	2 437	2 385	2 514	2 261	2 209	2 207	2 105	2 458
75 50	20 268	1 860	1 711	1 858	1 706	1 769	1 659	1 748	1 764	1 528	1 549	1 442	1 674
80 55	12 231	1 054	991	1 027	1 069	1 080	988	1 105	1 073	985	986	921	952
85 60	7 733	659	654	656	630	671	644	690	682	609	626	607	605
89 平成1年	5 064	476	363	432	417	400	429	431	436	442	427	411	400
90 2	4 664	447	385	430	407	381	378	376	399	361	377	365	358
早期新生児死亡数 Early neonatal deaths													
1970 昭和45年	12 810	1 098	1 023	1 110	1 025	1 031	1 043	1 185	1 080	1 011	1 049	1 011	1 144
75 50	10 245	873	812	847	832	887	886	1 012	864	802	817	775	838
80 55	6 154	490	481	519	491	540	526	528	531	532	463	510	543
85 60	3 737	302	298	327	286	372	325	310	325	297	327	254	314
89 平成1年	2 386	190	161	189	209	214	191	202	206	193	199	208	224
90 2	2 337	189	147	195	210	201	199	182	214	208	204	186	202
周産期死亡率 Perinatal death rates													
1970 昭和45年	21.7	22.1	22.9	23.5	21.2	21.6	22.4	21.8	20.4	20.5	21.3	20.9	21.5
75 50	16.0	16.3	16.5	16.9	15.9	16.5	16.7	16.4	16.0	14.6	15.3	15.5	16.0
80 55	11.7	11.4	11.8	11.9	12.2	12.1	11.8	11.8	11.6	11.1	10.9	11.9	11.7
85 60	8.0	8.0	8.6	8.5	8.0	8.6	8.3	7.7	8.0	7.4	7.7	7.7	7.7
89 平成1年	6.0	6.3	5.6	6.0	6.2	5.7	6.1	5.8	5.9	6.1	6.0	6.1	5.8
90 2	5.7	6.1	5.7	6.2	6.2	5.5	5.8	5.2	5.7	5.6	5.6	5.7	5.5
妊娠満28週以後の死産比 Foetal death ratio of 28 weeks and over of gestation													
1970 昭和45年	15.0	15.8	16.3	16.7	15.0	15.1	15.6	14.8	13.8	14.1	14.4	14.1	14.7
75 50	10.7	11.1	11.2	11.6	10.7	11.0	10.9	10.4	10.7	9.6	10.0	10.1	10.7
80 55	7.8	7.8	7.9	7.9	8.3	8.0	7.7	8.0	7.8	7.2	7.4	7.6	7.5
85 60	5.4	5.5	5.9	5.7	5.5	5.5	5.5	5.3	5.4	5.0	5.1	5.4	5.1
89 平成1年	4.1	4.5	3.8	4.2	4.1	3.7	4.2	4.0	4.0	4.3	4.1	4.0	3.7
90 2	3.8	4.3	4.2	4.3	4.1	3.6	3.8	3.5	3.7	3.5	3.7	3.8	3.5
早期新生児死亡率 Early neonatal death rates													
1970 昭和45年	6.6	6.3	6.6	6.8	6.2	6.4	6.8	7.0	6.6	6.4	6.9	6.8	6.8
75 50	5.4	5.2	5.3	5.3	5.2	5.5	5.8	6.0	5.3	5.0	5.3	5.4	5.3
80 55	3.9	3.6	3.8	4.0	3.8	4.0	4.1	3.8	3.8	3.9	3.5	4.2	4.3
85 60	2.6	2.5	2.7	2.8	2.5	3.1	2.8	2.4	2.6	2.4	2.7	2.3	2.6
89 平成1年	1.9	1.8	1.7	1.8	2.1	2.0	1.9	1.9	1.9	1.9	1.9	2.0	2.1
90 2	1.9	1.8	1.6	1.9	2.1	1.9	2.0	1.7	2.0	2.0	2.0	1.9	2.0

注: 1) 計算方法は53ページの「月間周産期死亡率」を参照されたい。

Perinatal death

[出産時の体重別にみた周産期死亡]

[Perinatal deaths by birth weight]

表 8.4 性・体重別にみた妊娠満28週以後の死産—早期新生児死亡別周産期死亡数
・率 (性・体重別出生千対) 及び百分率

Table 8.4 Perinatal deaths, perinatal death rates (per 1,000 live births)
and percent distribution by sex and birth weight: Japan, 1990

平成2年

	総数	~0.9 kg	1.0~1.4	1.5~1.9	2.0~2.4	2.5~2.9	3.0~3.4	3.5~3.9	4.0~	(再掲) 2.5 kg 以下 and under [Secondary [mention]
	Total									
周産期死亡数 ¹⁾ Perinatal deaths										
総数 Total	7 001	1 311	1 209	1 114	1 071	1 082	801	266	69	4 730
男 M.	3 717	663	643	583	549	563	462	169	47	2 450
女 F.	3 202	606	552	518	517	517	339	96	22	2 206
妊娠満28週以後の死産 Foetal deaths of 28 weeks and over of gestation	4 664	760	907	850	753	703	481	146	49	3 287
男 M.	2 394	364	473	436	374	359	260	88	34	1 654
女 F.	2 188	354	420	401	374	342	221	57	15	1 559
早期新生児死亡 Early neonatal deaths	2 337	551	302	264	318	379	320	120	20	1 443
男 M.	1 323	299	170	147	175	204	202	81	13	796
女 F.	1 014	252	132	117	143	175	118	39	7	647
周産期死亡率 ²⁾ Perinatal death rates										
総数 Total	5.7	572.2	286.0	103.1	17.8	2.9	1.4	1.4	3.0	59.6
男 M.	5.9	576.0	292.3	106.7	20.2	3.3	1.6	1.5	3.3	66.5
女 F.	5.4	531.6	272.3	97.1	15.7	2.5	1.3	1.3	2.6	52.0
妊娠満28週以後の死産比 Foetal death ratio of 28 weeks and over of gestation	3.8	331.7	214.6	78.7	12.5	1.9	0.9	0.8	2.1	41.4
男 M.	3.8	316.2	215.0	79.8	13.8	2.1	0.9	0.8	2.4	44.9
女 F.	3.7	310.5	207.2	75.1	11.4	1.7	0.8	0.8	1.8	36.7
早期新生児死亡率 Early neonatal death rates	1.9	240.5	71.4	24.4	5.3	1.0	0.6	0.6	0.9	18.2
男 M.	2.1	259.8	77.3	26.9	6.4	1.2	0.7	0.7	0.9	21.6
女 F.	1.7	221.1	65.1	21.9	4.4	0.9	0.4	0.5	0.8	15.2
百分率 ³⁾ Percentage										
総数 Total	100.0	18.9	17.5	16.1	15.5	15.6	11.6	3.8	1.0	68.3
男 M.	100.0	18.0	17.5	15.8	14.9	15.3	12.6	4.6	1.3	66.6
女 F.	100.0	19.1	17.4	16.4	16.3	16.3	10.7	3.0	0.7	69.7
妊娠満28週以後の死産 Foetal death of 28 weeks and over of gestation	100.0	16.3	19.5	18.3	16.2	15.1	10.3	3.1	1.1	70.7
男 M.	100.0	15.2	19.8	18.3	15.7	15.0	10.9	3.7	1.4	69.3
女 F.	100.0	16.2	19.2	18.4	17.1	15.7	10.1	2.6	0.7	71.4
早期新生児死亡 Early neonatal death	100.0	24.2	13.3	11.6	14.0	16.7	14.1	5.3	0.9	63.5
男 M.	100.0	23.2	13.2	11.4	13.6	15.8	15.6	6.3	1.0	61.7
女 F.	100.0	25.6	13.4	11.9	14.5	17.8	12.0	4.0	0.7	65.8

注: 1) 性・体重ともに不詳を含む。

2) 周産期死亡率 = $\frac{\text{年間の性・体重階級別周産期死亡数}}{\text{年間の性・体重階級別出生数}} \times 1,000$

3) 体重不詳を除き算出した。