

周産期死亡

〔月別にみた周産期死亡〕
〔Perinatal deaths by months〕

表 8.3 月別にみた妊娠満28週以後の死産—早期新生児死亡・
年次別周産期死亡数及び率¹⁾ (出生千対)

Table 8.3 Trends of perinatal deaths and perinatal death rates
(per 1,000 live births) by months : Japan

年 次	総 数	1 月	2 月	3 月	4 月	5 月	6 月	7 月	8 月	9 月	10 月	11 月	12 月
Year	Total	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
周産期死亡数 Perinatal deaths													
1970 昭和45年	41 917	3 864	3 545	3 856	3 522	3 468	3 428	3 699	3 341	3 220	3 256	3 116	3 602
75 50	30 513	2 733	2 523	2 705	2 538	2 656	2 545	2 760	2 628	2 330	2 366	2 217	2 512
80 55	18 385	1 544	1 472	1 546	1 560	1 620	1 514	1 633	1 604	1 517	1 449	1 431	1 495
85 60	11 470	961	952	983	916	1 043	969	1 000	1 007	906	953	861	919
88 63	8 508	760	656	692	704	731	747	754	747	655	696	690	676
89 平成1年	7 450	666	524	621	626	614	620	633	642	635	626	619	624
妊娠満28週以後の死産数 Foetal deaths of 28 weeks and over of gestation													
1970 昭和45年	29 107	2 766	2 522	2 746	2 497	2 437	2 385	2 514	2 261	2 209	2 207	2 105	2 458
75 50	20 268	1 860	1 711	1 858	1 706	1 769	1 659	1 748	1 764	1 528	1 549	1 442	1 674
80 55	12 231	1 054	991	1 027	1 069	1 080	988	1 105	1 073	985	986	921	952
85 60	7 733	659	654	656	630	671	644	690	682	609	626	607	605
88 63	5 759	525	453	478	471	501	506	510	498	435	454	467	461
89 平成1年	5 064	476	363	432	417	400	429	431	436	442	427	411	400
早期新生児死亡数 Early neonatal deaths													
1970 昭和45年	12 810	1 098	1 023	1 110	1 025	1 031	1 043	1 185	1 080	1 011	1 049	1 011	1 144
75 50	10 245	873	812	847	832	887	886	1 012	864	802	817	775	838
80 55	6 154	490	481	519	491	540	526	528	531	532	463	510	543
85 60	3 737	302	298	327	286	372	325	310	325	297	327	254	314
88 63	2 749	235	203	214	233	230	241	244	249	220	242	223	215
89 平成1年	2 386	190	161	189	209	214	191	202	206	193	199	208	224
周産期死亡率 Perinatal death rates													
1970 昭和45年	21.7	22.1	22.9	23.5	21.2	21.6	22.4	21.8	20.4	20.5	21.3	20.9	21.5
75 50	16.0	16.3	16.5	16.9	15.9	16.5	16.7	16.4	16.0	14.6	15.3	15.5	16.0
80 55	11.7	11.4	11.8	11.9	12.2	12.1	11.8	11.6	11.1	10.9	11.9	11.7	11.7
85 60	8.0	8.0	8.6	8.5	8.0	8.6	8.3	7.7	8.0	7.4	7.7	7.7	7.7
88 63	6.5	7.1	6.6	6.5	6.7	6.7	6.9	6.5	6.3	5.7	6.1	6.5	6.3
89 平成1年	6.0	6.3	5.6	6.0	6.2	5.7	6.1	5.8	5.9	6.1	6.0	6.1	5.8
妊娠満28週以後の死産比 Foetal death ratio of 28 weeks and over of gestation													
1970 昭和45年	15.0	15.8	16.3	16.7	15.0	15.1	15.6	14.8	13.8	14.1	14.4	14.1	14.7
75 50	10.7	11.1	11.2	11.6	10.7	11.0	10.9	10.4	10.7	9.6	10.0	10.1	10.7
80 55	7.8	7.8	7.9	7.9	8.3	8.0	7.7	8.0	7.8	7.2	7.4	7.6	7.5
85 60	5.4	5.5	5.9	5.7	5.5	5.5	5.5	5.3	5.4	5.0	5.1	5.4	5.1
88 63	4.4	4.9	4.6	4.5	4.5	4.6	4.7	4.4	4.2	3.8	4.0	4.4	4.3
89 平成1年	4.1	4.5	3.8	4.2	4.1	3.7	4.2	4.0	4.0	4.3	4.1	4.0	3.7
早期新生児死亡率 Early neonatal death rates													
1970 昭和45年	6.6	6.3	6.6	6.8	6.2	6.4	6.8	7.0	6.6	6.4	6.9	6.8	6.8
75 50	5.4	5.2	5.3	5.3	5.2	5.5	5.8	6.0	5.3	5.0	5.3	5.4	5.3
80 55	3.9	3.6	3.8	4.0	3.8	4.0	4.1	3.8	3.8	3.9	3.5	4.2	4.3
85 60	2.6	2.5	2.7	2.8	2.5	3.1	2.8	2.4	2.6	2.4	2.7	2.3	2.6
88 63	2.1	2.2	2.1	2.0	2.2	2.1	2.2	2.1	2.1	1.9	2.1	2.1	2.0
89 平成1年	1.9	1.8	1.7	1.8	2.1	2.0	1.9	1.9	1.9	1.9	1.9	2.0	2.1

注: 1) 計算方法は52ページの「月間周産期死亡率」を参照されたい。

Perinatal death

〔出産時の体重別にみた周産期死亡〕
〔Perinatal deaths by birth weight〕

表 8.4 性・体重別にみた妊娠満28週以後の死産—早期新生児死亡別周産期死亡数・率 (性・体重別出生千対) 及び百分率

Table 8.4 Perinatal deaths, perinatal death rates (per 1,000 live births) and percent distribution by sex and birth weight: Japan, 1989

平成元年

	総 数	~0.9 kg	1.0~1.4	1.5~1.9	2.0~2.4	2.5~2.9	3.0~3.4	3.5~3.9	4.0~	(再掲) 2.5 kg 以下 2.5 kg and under [Secondary mention]
	Total									
周産期死亡数 ¹⁾ Perinatal deaths										
総 数 Total	7 450	1 317	1 216	1 248	1 165	1 166	889	287	97	4 979
男 M.	3 880	654	665	655	575	584	483	172	61	2 564
女 F.	3 499	622	542	580	585	582	406	115	35	2 347
妊娠満28週以後の死産 Foetal death of 28 weeks and over of gestation	5 064	771	920	936	846	793	548	171	67	3 490
男 M.	2 577	361	497	478	415	391	292	96	42	1 760
女 F.	2 416	369	414	445	426	402	256	75	24	1 662
早期新生児死亡 Early neonatal deaths	2 386	546	296	312	319	373	341	116	30	1 489
男 M.	1 303	293	168	177	160	193	191	76	19	804
女 F.	1 083	253	128	135	159	180	150	40	11	685
周産期死亡率 ²⁾ Perinatal death rates										
総 数 Total	6.0	616.0	287.3	119.3	19.8	3.1	1.5	1.4	3.9	64.0
男 M.	6.1	630.1	300.6	124.4	21.5	3.4	1.6	1.5	3.9	70.9
女 F.	5.8	565.5	268.3	111.5	18.3	2.8	1.5	1.4	3.9	56.5
妊娠満28週以後の死産比 Foetal death ratio of 28 weeks and over of gestation	4.1	360.6	217.4	89.4	14.4	2.1	1.0	0.9	2.7	44.9
男 M.	4.0	347.8	224.7	90.8	15.5	2.3	1.0	0.8	2.7	48.6
女 F.	4.0	335.5	205.0	85.6	13.3	2.0	0.9	0.9	2.7	40.0
早期新生児死亡率 Early neonatal death rates	1.9	255.4	69.9	29.8	5.4	1.0	0.6	0.6	1.2	19.2
男 M.	2.0	282.3	75.9	33.6	6.0	1.1	0.6	0.6	1.2	22.2
女 F.	1.8	230.0	63.4	26.0	5.0	0.9	0.6	0.5	1.2	16.5
百 分 率 ³⁾ Percentage										
総 数 Total	100.0	17.8	16.5	16.9	15.8	15.8	12.0	3.9	1.3	67.4
男 M.	100.0	17.0	17.3	17.0	14.9	15.2	12.5	4.5	1.6	66.6
女 F.	100.0	17.9	15.6	16.7	16.9	16.8	11.7	3.3	1.0	67.7
妊娠満28週以後の死産 Foetal death of 28 weeks and over gestation	100.0	15.3	18.2	18.5	16.7	15.7	10.8	3.4	1.3	69.1
男 M.	100.0	14.0	19.3	18.6	16.1	15.2	11.4	3.7	1.6	68.4
女 F.	100.0	15.3	17.2	18.5	17.7	16.7	10.6	3.1	1.0	68.9
早期新生児死亡 Early neonatal deaths	100.0	23.4	12.7	13.4	13.7	16.0	14.6	5.0	1.3	63.8
男 M.	100.0	22.9	13.2	13.9	12.5	15.1	15.0	6.0	1.5	63.0
女 F.	100.0	24.0	12.1	12.8	15.1	17.0	14.2	3.8	1.0	64.9

注: 1) 性・体重ともに不詳を含む。

2) 周産期死亡率 = $\frac{\text{年間の性・体重階級別周産期死亡数}}{\text{年間の性・体重階級別出生数}} \times 1,000$

3) 体重不詳を除き算出した。